

1 THE SELF-GRAVITY MODEL OF LONGITUDINAL SPAN
2 OF NEPTUNE MAIN ARC FRATERNITE

3 K.H. Tsui and J.A. Souza

4 Instituto de Fisica - Universidade Federal Fluminense,
5 Campus da Praia Vermelha, Av. General Milton Tavares de Souza s/n,
6 Gragoata, 24.210-346, Niteroi, Rio de Janeiro, Brasil.

7 `tsui@if.uff.br`

8 Received _____; accepted _____

ABSTRACT

According to recent works [Tsui PSS **55**, 237-242 (2007); **55**, 2042-2044 (2007)], the Neptune Adams ring main arc Fraternite is regarded as captured by the corotation elliptic resonance (CER) potential of Galatea. The minor arcs Egalite(2,1), Liberte, and Courage are located at positions where the time averaged forces, due to the 42-43 corotation-Lindblad resonances under the central field of Neptune, vanish. With adequately chosen Fraternite mass and Galatea eccentricity, this model gives minor arc locations compatible to observed positions, and allows a dynamic transport of materials among arcs. To complement this model, the effect of self-gravity of Fraternite, with a distributed mass, is evaluated together with the CER potential to account for its 10^0 longitudinal span. Although self-gravity is the collective action of all the particles in the arc, each individual particle will see the self-potential with a central maximum as an external potential generated by other particles.

Subject headings: Planets: Rings

1. Introduction

From the very first observations of the Neptune Adams ring arcs [Hubbard et al 1986, Smith et al 1989] plus the subsequent observations [Sicardy et al 1999, Dumas et al 1999], the Adams arcs seemed to change in arc locations and in brightness. More recently, these dynamic natures of the arcs, Fraternite, Egalite (2,1), Liberte, and Courage, have been confirmed beyond any doubt in another ground observation [de Pater et al 2005]. Measuring from the center of the main arc Fraternite, they extend a total of about 40° ahead of Fraternite. Occassionally, some arcs flare up and others fade away. Furthermore, the arc configuration appears to be changing in time as well. The leading arc Courage appears to have leaped over to another CER site recently [de Pater et al 2005]. Although the twin arc Egalite(2,1) is small, it is a very bright arc. According to de Pater et al [2005], its relative intensity to Fraternite varied from 17 percent higher in 2002 to seven percent lower in 2003 totaling a 24 percent relative change over a short period of time. The angular span of the twin arc Egalite appeared to be 30 percent larger in 2005 and 1999 publications than in 1989 Voyager 2 results. This widening of Egalite was accompanied by a corresponding narrowing of Fraternite, which indicated a likely exchange of material between the two. As for Liberte, 1999 data showed it was about 3° ahead of its position in Voyager 2 pictures. For the 2005 results, the 2002 data appeared to show Liberte as a twin arc separated by about 4.5° with the leading twin at the original Voyager 1989 location, while in 2003 it returned again as one single arc at the Voyager location. With respect to the normally low intensity arc Courage, it flared in intensity to become as bright as Liberte in 1998 indicating a possible exchange of material between the two arcs. Most interestingly, it was observed in the 2005 data that Courage has moved 8° ahead from 31.2° to 39.7° [de Pater et al 2005].

According to the prevailing theories, based on the restricted three-body framework (Neptune-Galatea-arcs) with a conservative disturbing potential, these arcs are radially and

37 longitudinally confined by the corotation resonance potential of the inner moon Galatea.
38 In order to account for these arcs, the 84/86 corotation resonance due to the inclination
39 of Galatea (CIR) had been invoked to give a potential site of 4.18° [Goldreich et al 1986].
40 Later on the 42/43 resonance because of its eccentricity (CER) was considered giving a
41 resonant site of 8.37° on the Adams ring arcs [Porco 1991, Horanyi and Porco 1993, Foryta
42 and Sicardy 1996]. The arc particles librate about the potential maximum imposed by the
43 corotational resonance satellite Galatea. Dissipated energy of the particle is replenished by
44 the Lindblad resonance. Nevertheless, well established as it is, there are several difficulties.
45 Firstly, with Fraternite centered at the potential maximum spanning approximately 5° on
46 each side, it crosses two unstable potential points which ought to reduce the angular spread.
47 Secondly, the minor arcs leading ahead of Fraternite are mislocated with the CIR or CER
48 potential maxima. Furthermore, should the arcs were confined by the corotation potential,
49 there ought to be arcs in other locations along the Adams ring distributed randomly instead
50 of clustered near Fraternite.

2. Time-Dependent Arcs

Recently, there is a model that considers Fraternite as being captured by the CER potential of Galatea. With Fraternite having a finite mass, the minor arcs are clustered at locations along the Adams ring where the time averaged force vanishes under the corotation-Lindblad resonances [Tsui 2007a, Tsui 2007b]. The finite mass of Fraternite has been suggested by Namouni and Porco [2002] to pull on the pericenter precession of Galatea to account for the mismatch between the CER pattern speed and the mean motion of the arcs. The arc locations are determined by the Lindblad resonance reaction of the arc itself. Because the force vanishes only on a time averaged base, as comparing to the stationary CER potential in the rotating frame, the arc material could migrate on a long time scale from one site to another leading to flaring of some arcs and fading of others. This could also generate twin arcs (Egalite, Liberte) and displace Courage from 31.2° to 39.7° (resonant jump) [de Pater et al 2005], as required by observations. Although there are only arcs in the leading positions ahead, arcs in the trailing positions behind could be allowed in this model. According to this Lindblad reaction model, only Fraternite f is confined by the externally imposed CER potential of Galatea x which reads

$$\Phi_c = \frac{Gm_x}{a_x} \frac{1}{2} \left(2n + a_x \frac{\partial}{\partial a_x} \right) \frac{1}{a_x} b_{1/2}^{(n)}(\alpha) e_x \cos \phi_{fx} \quad , \quad (1)$$

where $\vec{r}_x = (r_x, \theta_x)$ and $\vec{r} = (r, \theta)$ are the position vectors of Galatea x and Fraternite mass distribution, a_x and a are the respective semi-major axes, ϕ_x is the argument of perihelion of x, $\phi_{fx} = (n\theta - (n-1)\theta_x - \phi_x)$ is the corotation resonance variable, $\alpha = a_x/a < 1$, and $n = 43$. With $a_x = 61952.60$ Km, $a = 62932.85$ Km, and $\alpha = 0.98444$ [Sicardy et al 1999, Dumas et al 1999], the CER potential is

$$\Phi_c = \frac{Gm_x}{a_x} 34e_x \cos \phi_{fx} . \quad (2)$$

72 To complement this model, we consider the self-gravity of Fraternite, which has a
 73 distributed mass, on the CER potential to account for its longitudinal 10^0 arc span. We
 74 first consider a qualitative spherical self-gravity physical model to grasp the 10^0 arc span.
 75 We begin with the Gauss law of the gravitational field

$$\nabla \cdot \vec{g}(\vec{r}) = -4\pi G\rho(\vec{r}) , \quad (3)$$

$$\vec{g} = +\nabla\Phi . \quad (4)$$

76 Under a qualitative physical model of arc span, we take a spherical uniform mass distribution
 77 of radius r_0 . Solving for the potential Φ inside the sphere with $\rho(\vec{r}) = \rho_0$ and outside the
 78 sphere with $\rho(\vec{r}) = 0$ respectively, and matching the potential and the gravitational field
 79 across the boundary, we get

$$\Phi_f = -\frac{1}{2} \frac{Gm_f}{r_0} \left(\frac{r_*}{r_0}\right)^2 + \frac{3}{2} \frac{Gm_f}{r_0} , \quad 0 < r_* < r_0 , \quad (5)$$

$$\Phi_f = +\frac{Gm_f}{r_*} , \quad r_0 < r_* < \infty . \quad (6)$$

80 This potential shows a normal $1/r_*$ decaying form for $r_0 < r_*$, but a r_*^2 form for $r_* < r_0$.
 81 Writing in terms of a_x and m_x , we have

$$\Phi_f = -\frac{1}{2} \frac{Gm_f a_x}{a_x r_0} \left(\frac{r_*}{r_0}\right)^2 + \frac{3}{2} \frac{Gm_f}{r_0} = -\frac{1}{2} \frac{Gm_x m_f a_x}{a_x m_x r_0} \left(\frac{a}{r_0}\right)^2 (\delta\theta)^2 + \frac{3}{2} \frac{Gm_x m_f a_x}{a_x m_x r_0} , \quad 0 < r_* < r_0 , \quad \delta\theta < \delta\theta_0 , \quad (7)$$

$$\Phi_f = +\frac{Gm_f a_x}{a_x r_*} = +\frac{Gm_x m_f}{a_x m_x} \frac{1}{\delta\theta} , \quad r_0 < r_* < \infty , \quad \delta\theta_0 < \delta\theta , \quad (8)$$

82 where r_* is now taken on the longitudinal direction along the arc, so that we can write
 83 $r_* = a\delta\theta$ and $r_0 = a\delta\theta_0$ with $\delta\theta$ as the angular span in radian. Taking $m_f/m_x = 10^{-3}$,
 84 $e_x = 10^{-4}$, and $\delta\theta_0 = 5^0 = 0.087\text{ rad}$, which are within the estimates of the arc parameters
 85 [Namouni and Porco 2002], we have plotted in Fig.1 the sinusoidal CER potential in
 86 thick line with a minimum around $\delta\theta = 4^0$ and the self-potential in thin line in units of
 87 Gm_x/a_x . The superposition of the two in thick line is also shown in the same figure. The
 88 superimposed potential has a maximum at the center and a minimum around $\delta\theta = 5^0$.
 89 Although self-gravity is resulted from all the particles of the arc, each individual particle
 90 will see the self-potential as an external potential. The particles will girate in stable orbit
 91 about the central maximum of the superpositioned CER potential and self-potential.

3. Self-Gravity

We now present an elongated ellipsoid model of self-gravity. For an ellipsoidal mass distribution with uniform density ρ_0 over a volume

$$\left(\frac{x}{a_1}\right)^2 + \left(\frac{y}{a_2}\right)^2 + \left(\frac{z}{a_3}\right)^2 = 1 \quad , \quad (9)$$

where $a_1 > a_2 > a_3$, the potential in space for the gravitational field $\vec{g}(\vec{r})$ had been addressed to in honorable treatises such as Kellogg [1953] and Landau and Lifshitz [1975]. Here, we follow the celebrated original work of Kellogg [1953] especially on Section 6 of Chapter 7. The potential in space of this homogeneous ellipsoid is given by

$$\Phi_f(x, y, z) = G\rho_0\pi a_1 a_2 a_3 \int_0^\infty \left[1 - \frac{x^2}{a_1^2 + \lambda} - \frac{y^2}{a_2^2 + \lambda} - \frac{z^2}{a_3^2 + \lambda}\right] \frac{d\lambda}{\sigma^{1/2}(\lambda)} \quad , \quad (10)$$

$$\sigma(\lambda) = (a_1^2 + \lambda)(a_2^2 + \lambda)(a_3^2 + \lambda) \quad , \quad (11)$$

where λ parameterizes a family of ellipsoids. Consider a prolate ellipsoid with $a_1 > a_2 = a_3$. This ellipsoid has a circular cross section on the yz plane and an axis of symmetry in x . The yz plane of $x = 0$ is the equatorial plane. In this prolate case, the self-potential inside and outside the ellipsoid is given respectively by [Exercise 6, p.196 of Kellogg]

$$\begin{aligned} \Phi_f(x, r) = G\rho_0 \frac{4\pi}{3} a_1 a_2 a_3 \frac{1}{f^2} & \left[(4x^2 - 2r^2 - f^2) \frac{1}{2f} \ln\left(\frac{2a_1 - f}{2a_1 + f}\right)^{1/2} \right. \\ & \left. + \frac{4a_1^2(2x^2 - r^2) - 2f^2x^2}{2a_1(4a_1^2 - f^2)} \right] \quad , \end{aligned} \quad (12)$$

$$\begin{aligned} \Phi_f(x, r) = G\rho_0 \frac{4\pi}{3} a_1 a_2 a_3 \frac{1}{f^2} & \left[(4x^2 - 2r^2 - f^2) \frac{1}{2f} \ln\left(\frac{s - f}{s + f}\right)^{1/2} \right. \\ & \left. + \frac{s^2(2x^2 - r^2) - 2f^2x^2}{s(s^2 - f^2)} \right] \quad , \end{aligned} \quad (13)$$

where f is the distance between the two foci with $(f/2)^2 = a_1^2 - a_2^2$, r is the perpendicular distance to the axis of symmetry with $r^2 = y^2 + z^2$, s is the sum of distances from the two foci to the point of interest $\vec{r}$. The inside potential can be obtained from the outside potential by using $s = 2a_1$. To evaluate the potential on the axis of symmetry, we take $r = 0$. Denoting $m_f = \rho_0(4\pi/3)a_1a_2a_3$ and considering $a_1 \gg a_2$, we get

$$\begin{aligned} \Phi_f(x) = & -\frac{Gm_f}{a_x} \frac{a_x}{4a_1} [\ln(\frac{2a_1}{a_2}) - 1] (\frac{\delta\theta}{\delta\theta_0})^2 \\ & + \frac{Gm_f}{a_x} \frac{a_x}{4a_1} \ln(\frac{2a_1}{a_2}) , \quad \delta\theta < \delta\theta_0 , \end{aligned} \quad (14)$$

$$\begin{aligned} \Phi_f(x) = & -\frac{Gm_f}{a_x} \frac{a_x}{4a_1} [\frac{1}{2} \ln(\frac{x+f/2}{x-f/2}) (\frac{\delta\theta}{\delta\theta_0})^2 - (\frac{\delta\theta}{\delta\theta_0})] \\ & + \frac{Gm_f}{a_x} \frac{a_x}{4a_1} \frac{1}{2} \ln(\frac{x+f/2}{x-f/2}) , \quad \delta\theta_0 < \delta\theta , \end{aligned} \quad (15)$$

for the self-potential inside and outside the ellipsoid respectively. Taking again $m_f/m_x = 10^{-3}$, with $a_x = 61952.60 \text{ Km}$ for Galatea, and semi-major axes $a_1 = 5500 \text{ Km}$ and $a_2 = 55 \text{ Km}$, the CER potential, the self-potential, and the superposition of the two with a minimum around $\delta\theta = 5^\circ$ are shown in Fig.2.

With Fraternite having 1×10^{-3} of the mass of Galatea, the self-potential actually exceeds the CER potential in magnitude, as shown in Fig.2. Each test mass would be librating around the potential maximum, dominated by the self-gravity of the collective mass distribution. Should Fraternite be elongated further while maintaining the total mass, it would increase the semi-major axis a_1 of the ellipsoid. This would reduce the amplitude of the self-potential of Eq. 14 through the $(a_x/4a_1)$ factor in the constant term, and weaken the self-potential. The elongation would feed the minor arcs. With this self-gravity model, not just the minor arcs are dynamically changing [de Pater et al 2005], the main arc Fraternite could be under a dynamical process as well.

4. Conclusions

In order to explain the 10^0 arc span of Fraternite, we draw attention to the fact that Fraternite, as an arc, has a significant mass. This mass is a distributed mass, instead of a point-like mass, such that its self-gravity should be taken into considerations to account for its angular span. We have used two models to evaluate the self-potential in the longitudinal direction. First is the tutorial spherical model, as a proof of principle study, with a uniform mass distribution over a sphere of radius r_0 . Second is the elongated ellipsoidal model for a more realistic evaluation. Using the accepted range of Fraternite parameters, the ellipsoid model shows that the self-potential of the arc could be the cause of its angular span. For a longer arc, the ellipsoid gets longer and the ratio a_1/a_2 becomes larger. Eventually, for a complete ring, the ellipsoid is infinitely long and the self-potential in the longitudinal direction becomes constant. The effects of self-gravity are felt only in the transverse direction for a planetary ring.

REFERENCES

134

- 135 de Pater, I., et al 2005. "The dynamic neptunian ring arcs: evidence for a gradual
136 disappearance of Liberte and resonant jump of Courage". Icarus 174, 263-272.
- 137 Dumas, C., et al 1999. "Stability of Neptune's ring arcs in question". Nature 400, 733-735.
- 138 Foryta, D.W., and B. Sicardy 1996. "The dynamics of the Neptunian Adams ring's arcs".
139 Icarus 123, 129-167.
- 140 Goldreich, P., S. Tremaine, and N. Borderies 1986. "Towards a theory for Neptune's arc
141 rings". Astron. J. 92, 490-494.
- 142 Horanyi, M., and C.C. Porco 1993. "Where exactly are the arcs of Neptune?". Icarus 106,
143 525-535.
- 144 Hubbard, W.B., et al 1986. "Occultation detection of a neptunian ring like arc". Nature
145 319, 636-640.
- 146 Kellogg, O.D. 1953. "Foundations of Potential Theory". Dover Publications, New York.
- 147 Landau, L.D., and E.M. Lifshitz 1975. "The Classical Theory of Fields". Pergamon Press,
148 Oxford.
- 149 Namouni, F., and C.C. Porco 2002. "The confinement of Neptune's ring arcs by the moon
150 Galatea". Nature 417, 45-47.
- 151 Porco, C.C. 1991. "An explanation for Neptune's ring arcs". Science 253, 995-1001.
- 152 Sicardy, B., et al 1999. "Images of Neptune's ring arcs obtained by a ground based
153 telescope". Nature 400, 731-732.
- 154 Smith, B.A., et al 1989. "Voyager 2 at Neptune: imaging science results". Science 246,
155 1422-1449.

- 156 Tsui, K.H. 2007a. "The configuration of Fraternite-Egalite2-Egalite1 in the Neptune ring
157 arcs system". Planetary Space Science 55, 237-242.
- 158 Tsui, K.H. 2007b. "The dynamic nature of the Adams ring arcs - Fraternite, Egalite(2,1),
159 Liberte, Courage". Planetary Space Science 55, 2042-2044.




Fig. 1.— The CER sinusoidal potential of Galatea in thick line, the self-potential of Fraternite with spherical model in thin line, and the sum of the two in thick line are plotted in units of Gm_x/a_x .




Fig. 2.— The CER sinusoidal potential of Galatea in thick line, the self-potential of Fraternite with ellipsoidal model in thin line, and the sum of the two in thick line are plotted in units of Gm_x/a_x .